

## Some remarks on multilinear maps and interpolation

Loukas Grafakos · Nigel Kalton

Received January 31, 1999 / Accepted February 27, 2000 /

Published online December 8, 2000 – © Springer-Verlag 2000

**Abstract.** A multilinear version of the Boyd interpolation theorem is proved in the context of quasi-normed rearrangement-invariant spaces. A multilinear Marcinkiewicz interpolation theorem is obtained as a corollary. Several applications are given, including estimates for bilinear fractional integrals.

*Mathematics Subject Classification (1991):* 46B70, 46E30, 42B99

### 1. Introduction

In this article we give a version of the Boyd interpolation theorem for multilinear operators. We will be working with rearrangement invariant quasi-Banach spaces, which include all the well-known examples such as Orlicz spaces and Lorentz spaces.

We will consider the following situation. Consider  $\mathbf{R}_+ = (0, \infty)$  with Lebesgue measure (which can of course be replaced by any infinite nonatomic measure space). We let  $L_0(0, \infty)$  be the space of all real-valued measurable functions equipped with the topology of local convergence in measure. Let  $\mathcal{E}$  be the space of all measurable functions which are bounded and supported on sets of finite measure. Now let  $T : \mathcal{E}^n \rightarrow L_0(0, \infty)$  be a multilinear map (our results also apply to sublinear maps). We suppose that  $T$  is locally continuous i.e. continuous when restricted to  $\prod_{k=1}^n L_\infty(E_k)$  for every choice of sets  $E_k$  of finite measure. We also suppose that  $T$  obeys a finite collection of weak type inequalities

$$\|T(\chi_{E_1}, \dots, \chi_{E_n})\|_{L_{p,\infty}} \leq C \prod_{k=1}^n |E_k|^{\theta_k}$$

for every  $n$ -tuple of measurable sets  $(E_1, \dots, E_n)$ . Here  $L_{p,\infty}$  is the usual weak  $L_p$  space and  $\theta_k > 0$  for every  $k$ . We then seek to characterize  $(n + 1)$ -tuples

---

L. GRAFAKOS, N. KALTON

Department of Mathematics, University of Missouri, Columbia, MO 65211, USA

(e-mail: {loukas,nigel}@math.missouri.edu)

Research of both authors was partially supported by the NSF

of rearrangement-invariant spaces  $(X_1, \dots, X_n, Y)$  for which  $T$  extends to a bounded  $n$ -linear map from  $X_1 \times \dots \times X_n$  into  $Y$ . In general one needs two distinct hypotheses. The first consists of an assumption on the Boyd indices of the spaces  $X_1, \dots, X_n, Y$ , as in the original Boyd interpolation theorem. The second hypothesis is that a certain  $n$ -linear test map associated with  $T$  is continuous.

Our main result (Theorem 4.1) gives a necessary (and often sufficient) condition on  $(X_1, \dots, X_n, Y)$  in the case when one has  $n+1$  such conditions which are sufficiently independent. Note that the original theorem of Boyd [2] corresponds to the case when  $n = 1$  and there are two conditions of the type:

$$\|T(\chi_E)\|_{L_{p,\infty}} \leq C|E|^{1/p}.$$

We deduce Theorem 4.1 from a similar *homogeneous* Boyd-type theorem (Theorem 3.7) which is applicable for example to  $n$ -linear forms. As a corollary we obtain a multilinear version of the Marcinkiewicz interpolation theorem (Theorem 4.6).

Our work is related to work of Strichartz [17], Janson [5], and Christ [3]. Note that as in [5] and [17] (and in contrast to [3]) our multilinear assumptions consist only of a finite number of estimates. Our results also develop and extend earlier work of Sharpley (see [15], [16], and [1]).

In Sect. 5 we give examples of multilinear interpolation. As one of our applications, we characterize the indices  $(1/p, 1/q, 1/r)$ ,  $0 < p, q, r \leq \infty$ , for which the bilinear fractional integral operator

$$I_\alpha(f, g)(x) = \int_{\mathbf{R}^n} f(x+t)g(x-t)|t|^{\alpha-n} dt.$$

maps  $L_p(\mathbf{R}^n) \times L_q(\mathbf{R}^n) \rightarrow L_r(\mathbf{R}^n)$ . This characterization was also independently obtained by C. Kenig and E. M. Stein [10].

## 2. Preliminaries

In this section we set up the background required to state the multilinear Boyd interpolation theorem.

Let  $L_0(0, \infty)$  be the space of all complex-valued measurable functions on  $(0, \infty)$ , with the topology of local convergence in measure. We define a quasi-Banach function space  $X$  on  $(0, \infty)$  to be a subspace of  $L_0$  equipped with a quasi-norm  $\|\cdot\|_X$  such that:

- $\|f\|_X = 0$  if and only if  $f = 0$  a.e.
- $\|\alpha f\|_X = |\alpha| \|f\|_X$ , whenever  $f \in X$  and  $\alpha \in \mathbf{C}$ .
- There exists a constant  $C$  so that if  $f, g \in X$  then  $\|f + g\|_X \leq C(\|f\|_X + \|g\|_X)$ .
- $X$  is complete (i.e. a quasi-Banach space) for  $\|\cdot\|_X$ .

- The injection  $X \rightarrow L_0$  is continuous.
- If  $E$  is a set of finite measure then  $\chi_E \in X$ .
- If  $f \in X$  and  $g \in L_0$  with  $|g| \leq |f|$  a.e. then  $g \in X$  and  $\|g\|_X \leq \|f\|_X$ .
- If  $0 \leq f_n \uparrow f$  a.e. and  $f \in X$  then  $\|f_n\|_X \uparrow \|f\|_X$ .

By assumption  $X$  must contain the space  $\mathcal{E}$  of all bounded measurable functions supported on sets of finite measure. We say that  $X$  is *minimal* if  $\mathcal{E}$  is dense in  $X$ . We say that  $X$  is *maximal* if it has the property that if  $0 \leq f_n \uparrow f$  a.e. with  $\sup \|f_n\|_X < \infty$ , then  $f \in X$ .

A quasi-Banach function space on  $(0, \infty)$  which is either maximal or minimal (cf. [12]) is said to be a *rearrangement-invariant function space* or *r.i. space* if  $\|f^*\|_X = \|f\|_X$  for all  $f \in X$ , where  $f^*$  is the decreasing rearrangement of  $|f|$ , i.e.  $f^*(t) = \inf\{x : |\{|f| > x\}| \leq t\}$ .

We say that  $X$  is  $r$ -convex if there is a constant  $C$  so that if  $f_1, \dots, f_n \in X$  then

$$\left\| \left( \sum_{i=1}^n |f_i|^r \right)^{1/r} \right\|_X \leq C \left( \sum_{i=1}^n \|f_i\|_X^r \right)^{1/r}.$$

For a discussion of  $r$ -convexity in the context of Banach lattices we refer to [12]; we refer to [7] for quasi-Banach lattices. Every Banach r.i. space is of course 1-convex, but there are examples of quasi-Banach r.i. spaces which fail to be  $r$ -convex for any  $r > 0$ , see [6]. However it is very natural to assume  $r$ -convexity since all “practical spaces” are  $r$ -convex for some  $r > 0$ .

Once an r.i. space  $X$  is defined on  $(0, \infty)$  it may be transferred to any  $\sigma$ -finite measure space  $(\Omega, \mu)$  by defining  $X(\Omega, \mu)$  to be the space of all measurable  $f : \Omega \rightarrow \mathbf{C}$  such that  $\|f\|_{X(\Omega)} = \|f^*\|_{X(0, \infty)} < \infty$ . In general if  $\Omega$  is a Polish space and  $\mu$  is an infinite nonatomic Borel measure there is a measure-preserving bijection of  $\Omega$  onto  $(0, \infty)$ . Thus there is no loss of generality in treating only the case of  $\Omega = (0, \infty)$ .

If  $X$  is an r.i. space then the dilation operators  $D_a : X \rightarrow X$  given by

$$(D_a f)(x) = f(x/a)$$

are well-defined and bounded. We define the Boyd indices by

$$p_X = \lim_{a \rightarrow \infty} \frac{\log a}{\log \|D_a\|}$$

and

$$q_X = \lim_{a \rightarrow 0} \frac{\log a}{\log \|D_a\|}.$$

Then  $0 < p_X \leq q_X \leq \infty$ . We refer to [12] or [1] for relevant discussion. If  $\epsilon > 0$  then there is a constant  $C = C(\epsilon, X)$  so that for all  $f \in X$  we have

$$(1) \quad \|D_a f\|_X \leq C \max \left( a^{\frac{1}{p_X} + \epsilon}, a^{\frac{1}{q_X} - \epsilon} \right).$$

It is sometimes useful to have the notion of a *carrier space* for an r.i. space  $X$ . Let  $\tilde{X}$  be a maximal quasi-Banach function space on  $(0, \infty)$  with the property that the dilation operators  $D_a$  are bounded on  $\tilde{X}$  and  $\|D_a\|_{\tilde{X}} \leq Ca^\kappa$  for some  $\kappa > 0$  and all  $a \geq 1$ . Then we can define an r.i. space  $X$  by requiring  $f \in X$  if and only if  $f^* \in \tilde{X}$  and by setting  $\|f\|_X = \|f^*\|_{\tilde{X}}$ . It is then easy to show that  $X$  is a maximal r.i. space and that  $\alpha_X \leq \kappa$ . We will in this case refer to  $\tilde{X}$  as a carrier space for  $X$ . Notice, of course, that  $X$  is a carrier space for itself.

Examples of r.i. spaces are provided by the usual Lorentz spaces  $L_{p,q}$  with (quasi)-norm

$$(2) \quad \|f\|_{L_{p,q}} = \begin{cases} \left( \int_0^\infty [f^*(t)t^{1/p}]^q \frac{dt}{t} \right)^{1/q} & \text{when } 0 < q < \infty, \\ \sup_{t>0} f^*(t)t^{1/p} & \text{when } q = \infty \end{cases}$$

for  $0 < p, q \leq \infty$ . These spaces are 1-convex (i.e. normable) when  $1 < p < \infty$  and  $1 \leq q \leq \infty$  or if  $p = q = 1$ . In general  $L_{p,q}$  is  $q$ -convex if  $q \leq p$  and  $s$ -convex for any  $s < p$  if  $q > p$ . The Boyd indices of  $L_{p,q}$  both coincide with  $p$ . Note that all these spaces have natural carrier spaces which are weighted  $L_p$ -spaces.

The significance of the Boyd indices lies in the fact that they can be used to characterize all rearrangement-invariant Banach spaces  $X$  on which certain known operators are bounded. For instance the Hardy-Littlewood maximal operator is bounded on  $X$  (r.i. over  $\mathbf{R}^n$ ) if and only if  $q_X < \infty$ , see [13], [18]. The Hilbert transform is bounded on  $X$  (r.i. over  $\mathbf{R}$ ) if and only if  $1 < p_X \leq q_X < \infty$ , see [2].

Let us now recall the Boyd interpolation theorem for  $(0, \infty)$  (see [2] or [12], p.145):

**Theorem 2.1.** *Suppose  $1 \leq p < q < \infty$  and that  $T : L_{p,1} + L_{q,1} \rightarrow L_0(0, \infty)$  is a linear map of weak types  $(p, p)$  and  $(q, q)$ . Suppose  $X$  is an r.i. space with  $p < p_X \leq q_X < q$ . Then  $T$  is a bounded map from  $X$  into itself.*

This result was extended to the case  $0 < p < q < \infty$  in [8] (Theorem 1.3) with the additional assumption that  $X$  is  $r$ -convex for some  $r > 0$ .

The main purpose of this article is to obtain a multilinear version of Theorem 2.1. This is achieved in the next two sections. We first obtain a homogeneous multilinear version of Theorem 2.1 (Theorem 3.7), and from this we deduce an inhomogeneous version, Theorem 4.1.

### 3. The homogeneous multilinear Boyd theorem

Let  $\mathcal{E}$  be the space of all measurable functions on  $(0, \infty)$  which are bounded and have support of finite measure. We shall say that a map (usually  $n$ -linear)

$T : \mathcal{E}^n \rightarrow Y$  in any topological vector space is locally continuous if its restriction to  $\prod_{k=1}^n L_\infty(E_k)$  is continuous for every choice of sets  $E_k$  of finite measure.

Now suppose  $\Theta$  is a finite subset of  $(\overline{\mathbf{R}}_+)^n = [0, \infty)^n$  and  $Y$  is a quasi-Banach space. We say that an  $n$ -linear map  $T : \mathcal{E}^n \rightarrow Y$  is  $\Theta$ -admissible if  $T$  is locally continuous and there is a constant  $M$  so that for every  $\theta = (\theta_1, \dots, \theta_k) \in \Theta$  we have

$$(3) \quad \|T(\chi_{E_1}, \dots, \chi_{E_n})\|_Y \leq M \prod_{k=1}^n |E_k|^{\theta_k},$$

whenever  $E_1, \dots, E_n$  have finite measure. The least such constant  $M$  is denoted by  $\|T\|_\Theta$ . In most of the work that follows, it will be convenient to take  $\Theta \subset \mathbf{R}_+^n$  i.e. to require  $\theta_k > 0$  for all  $\theta, k$ .

Let us recall that a quasi-Banach space  $(Y, \|\cdot\|)$  is called  $s$ -normed if there is a constant  $C$  such that for all  $y_1, \dots, y_m \in Y$  we have

$$\|y_1 + \dots + y_m\|^s \leq C(\|y_1\|^s + \dots + \|y_m\|^s).$$

Now let  $\mathbf{X} = (X_1, \dots, X_n)$  be an  $n$ -tuple of r.i. spaces. We say that  $\mathbf{X}$  has the *interpolation condition*  $(\Theta, s)$ , where  $0 < s \leq 1$ , if for every  $s$ -normed quasi-Banach space  $Y$  and every  $\Theta$ -admissible  $T : \mathcal{E}^n \rightarrow Y$  there is a continuous extension  $T : X_1 \times \dots \times X_n \rightarrow Y$  with norm a constant multiple of  $\|T\|_\Theta$ . Note here that in the case  $s = 1$  it is sufficient to take  $Y$  to be the scalar field  $\mathbf{R}$  or  $\mathbf{C}$  and hence we only consider  $n$ -linear forms.

We will need to establish some examples of  $\Theta$ -admissible multilinear maps. We begin with a lemma.

**Lemma 3.1.** *Suppose  $0 < u < \infty$  and  $0 < s < r < \infty$ . Then for any measurable set  $E \subset (0, \infty)$  we have*

$$\left( su \int_E x^{su-1} dx \right)^{1/s} \leq \left( ru \int_E x^{ru-1} dx \right)^{1/r}.$$

*In particular if  $su < 1$  then*

$$\left( su \int_E x^{su-1} dx \right)^{1/s} \leq |E|^u.$$

*Proof.* First note that for  $t > 1$ , we have  $(t^{su} - 1)^{1/s} \leq (t^{ru} - 1)^{1/r}$  and also that  $(t^{su} - 1)^{1/s} (t^{ru} - 1)^{-1/r}$  is increasing. This last fact follows from the observation that  $t \rightarrow \frac{1}{r} \log(t^r - 1) - \frac{1}{s} \log(t^s - 1)$  is monotone decreasing and converges to zero at infinity. This implies that if  $E$  is an interval we have the desired inequality.

We now proceed to prove the result for  $E$  a disjoint union of  $m$  intervals using induction. Assume the required inequality is true for all unions of less than  $m$  disjoint intervals. Now if  $E$  is a finite union of  $m$  disjoint intervals  $[v_j, w_j)$  for

$1 \leq j \leq m$  where  $v_1 < w_1 < \dots < v_m < w_m$ , we define  $h > w_{m-1}$  by the condition that

$$h^{su} - w_{m-1}^{su} = (w_m^{su} - v_m^{su}) + (w_{m-1}^{su} - v_{m-1}^{su}).$$

If we had

$$(4) \quad h^{ru} - w_{m-1}^{ru} \leq (w_m^{ru} - v_m^{ru}) + (w_{m-1}^{ru} - v_{m-1}^{ru}),$$

then the inductive hypothesis applied to the  $m - 1$  intervals  $[v_1, w_1), \dots, [v_{m-2}, w_{m-2})$ , and  $[v_{m-1}, h)$  together with (4) would quickly give the desired conclusion. It suffices therefore to prove (4). This will follow from the fact that if  $\alpha, \beta, \gamma$ , and  $\delta$  are positive numbers satisfying  $\alpha + \gamma = \beta + \delta$  and  $\beta < \gamma < \delta$ , then  $\alpha^{r/s} + \gamma^{r/s} \leq \beta^{r/s} + \delta^{r/s}$  when  $r > s$ . Indeed, the assumptions above imply that  $\beta < \alpha < \delta$  and clearly

$$\alpha^{r/s} + \gamma^{r/s} \leq \max_{\alpha \in (\beta, \delta)} (\alpha^{r/s} + (\beta + \delta - \alpha)^{r/s}) \leq \beta^{r/s} + \delta^{r/s}.$$

□

Let  $\langle \cdot, \cdot \rangle$  denote the usual inner product on  $\mathbf{R}^n$  and  $\| \cdot \|$  the usual Euclidean norm. For each  $\theta \in \mathbf{R}^n$  let  $\theta_k$  denote its  $k^{\text{th}}$  coordinate. Suppose  $\Theta$  is a finite subset of  $(\overline{\mathbf{R}}_+)^n = [0, \infty)^n$ . Define a sublinear map associated with  $\Theta$  as follows

$$(5) \quad a(\xi) = a_\Theta(\xi) = \max_{\theta \in \Theta} \langle \xi, \theta \rangle.$$

Let  $\mathbf{X} = (X_1, \dots, X_n)$  be an  $n$ -tuple of r.i. spaces. We have the following theorem. See also Sharpley [15] for a somewhat similar result.

**Theorem 3.2.** *Let  $0 < s \leq 1$ . Consider the statements:*

- (i)  $\mathbf{X}$  satisfies the interpolation condition  $(\Theta, s)$ .
- (ii) There exists a constant  $C$  so that if  $f_1, \dots, f_n \in \mathcal{E}$ ,

$$(6) \quad \left( \int_{\mathbf{R}^n} \prod_{k=1}^n (f_k^*(e^{\xi_k}))^s \exp(-sa(-\xi)) d\xi \right)^{1/s} \leq C \prod_{k=1}^n \|f_k\|_{X_k}.$$

- (iii) There exists a constant  $C$  so that if  $f_1, \dots, f_n \in \mathcal{E}$ ,

$$(7) \quad \left( \int_{\mathbf{R}^n} \prod_{k=1}^n (F_k(e^{\xi_k}))^s \exp(-sa(-\xi)) d\xi \right)^{1/s} \leq C \prod_{k=1}^n \|f_k\|_{X_k},$$

where

$$F_k(x) = \left( \frac{1}{x} \int_0^x (f_k^*(t))^s dt \right)^{1/s}.$$

(iv) There exists a constant  $C$  so that if  $f_1, \dots, f_n \in \mathcal{E}$ , then

$$(8) \quad \max_{\|\xi\|=1} \left( \int_0^\infty \prod_{k=1}^n (f_k^*(e^{t\xi_k}))^s \exp(-sta(-\xi)) dt \right)^{1/s} \leq C \prod_{k=1}^n \|f_k\|_{X_k}.$$

Then (ii) implies (i). Furthermore, if  $\Theta \subset \mathbf{R}_+^n$  and  $s$  is small enough so that  $s\theta_k < 1$  for every  $(\theta_1, \dots, \theta_k) \in \Theta$  and every  $1 \leq k \leq n$ , then (i), (ii), (iii), and (iv) are all equivalent.

*Proof.* First assume (ii) and that  $T : \mathcal{E}^n \rightarrow Y$  is  $\Theta$ -admissible where  $Y$  is  $s$ -normed. Without loss of generality we assume  $\|T\|_\Theta \leq 1$ . We first note that if  $f_k$  are supported in  $E_k$  and  $\|f_k\|_{L_\infty} \leq 1$  then we have an estimate:

$$(9) \quad \|T(f_1, \dots, f_n)\|_Y \leq C \min_{\theta \in \Theta} \prod_{k=1}^n |E_k|^{\theta_k},$$

where  $C$  depends only on  $s$  and  $n$ . To see this it suffices to get an estimate for positive functions  $f_k$  and then extend to signed and complex functions by additivity. But if  $f_k$  is positive we can write

$$f_k = \sum_{j=1}^\infty 2^{-j} \chi_{A_{jk}}$$

where  $A_{jk} \subset E_k$ . Expanding out we easily get estimate (9).

Now suppose  $f_1, \dots, f_n \in \mathcal{E}$ . We can write each  $f_k$  in the form

$$f_k = \sum_{m=-\infty}^\infty f_k \chi_{A_{km}}$$

where  $|A_{km}| = 2^m$  and  $\|f_k \chi_{A_{km}}\|_{L_\infty} \leq f_k^*(2^m)$ . Now by (9) we have

$$\|T(f_1 \chi_{A_{1m_1}}, \dots, f_n \chi_{A_{nm_n}})\|_Y \leq C \min_{\theta \in \Theta} 2^{\sum_{k=1}^n \theta_k m_k} \prod_{k=1}^n f_k^*(2^{m_k}).$$

Now since  $\|\cdot\|_Y$  is an  $s$ -norm after summing and making an obvious integral estimate we obtain

$$\|T(f_1, \dots, f_n)\|_Y^s \leq C \int_0^\infty \dots \int_0^\infty (f_1^*(x_1))^s \dots (f_n^*(x_n))^s \min_{\theta \in \Theta} \prod_{k=1}^n x_k^{s\theta_k - 1} dx_1 \dots dx_n.$$

The right-hand side is now estimated by  $C \prod_{k=1}^n \|f_k\|_{X_k}$  using (6). We can now extend the definition of  $T(f_1, \dots, f_n)$  to  $X_1 \times \dots \times X_n$  by noting that for  $f_k \in X_k$  the sum

$$\sum_{m_1, \dots, m_n} T(f_1 \chi_{A_{1m_1}}, \dots, f_n \chi_{A_{nm_n}})$$

converges in  $Y$ . It is easy to check that this extends  $T$  unambiguously and continuously to  $X_1 \times \dots \times X_n$ . Thus (i) holds.

Now assume (i),  $\Theta \subset \mathbf{R}_+^n$ , and  $s\theta_k < 1$  for all  $\theta \in \Theta$  and  $1 \leq k \leq n$ . For each  $\theta \in \Theta$  Lemma 3.1 gives that if  $f_k = \chi_E \in \mathcal{E}$ , then

$$\left( \int_0^\infty x^{s\theta_k-1} f_k(x)^s dx \right)^{1/s} \leq (s\theta_k)^{-1/s} |E|^{\theta_k},$$

where as usually  $\theta_k$  denotes the  $k^{\text{th}}$  coordinate of  $\theta$ . It follows that if we define

$$T_\theta(f_1, \dots, f_n)(x_1, \dots, x_n) = \prod_{k=1}^n x_k^{\theta_k - \frac{1}{s}} f_k(x_k),$$

then  $T_\theta : \mathcal{E}^n \rightarrow L_s((0, \infty)^n)$  is  $\{\theta\}$ -admissible and  $\|T_\theta\|_{\{\theta\}} \leq s^{-n/s} \prod_{k=1}^n \theta_k^{-1/s}$ . If we define  $T$  by

$$T(f_1, \dots, f_n)(x_1, \dots, x_n) = \prod_{k=1}^n \left( \min_{\theta \in \Theta} x_k^{\theta_k - \frac{1}{s}} \right) f_k(x_k)$$

then  $T$  is  $\Theta$ -admissible. It follows that we can find  $C$  so that (6) is valid and thus (ii) holds.

We now show that (ii) implies (iii). Observe that

$$(F_k(e^{\xi_k}))^s = \int_{-\infty}^{\xi_k} e^{\eta_k - \xi_k} (f_k^*(e^{\eta_k}))^s d\eta_k.$$

Hence

$$\begin{aligned} & \int_{\mathbf{R}^n} \prod_{k=1}^n (F_k(e^{\xi_k}))^s \exp(-sa(-\xi)) d\xi \\ (10) \quad & = \int_{\mathbf{R}^n} \int_{\eta \leq \xi} \prod_{k=1}^n (f_k^*(e^{\eta_k}))^s \exp(\langle \eta - \xi, \mathbf{1} \rangle - sa(-\xi)) d\eta d\xi, \end{aligned}$$

where  $\eta \leq \xi$  means  $\eta_k \leq \xi_k$  for  $1 \leq k \leq n$  and  $\mathbf{1}$  denotes the vector  $(1, 1, \dots, 1)$ . For fixed  $\eta$  pick  $\tilde{\theta} \in \Theta$  so that  $a(-\eta) = \langle \tilde{\theta}, -\eta \rangle$ . Then

$$\begin{aligned} \int_{\xi \geq \eta} \exp(-\langle \xi, \mathbf{1} \rangle - sa(-\xi)) d\xi & \leq \int_{\xi \geq \eta} \exp(-\langle \xi, \mathbf{1} - s\tilde{\theta} \rangle) d\xi \\ & \leq C \exp(-\langle \eta, \mathbf{1} \rangle - sa(-\eta)) \end{aligned}$$



since for some  $\delta > 0$  we have  $1 - s\theta_k > \delta > 0$  for all  $\theta \in \Theta$  and all  $k$ . Substituting back into (10) gives the required estimate.

Next we show (iii) implies (iv). To do this it suffices to note that the map  $\xi \rightarrow \sum_{k=1}^n \log F_k(e^{\xi k})$  is Lipschitz with a constant depending only on  $s$  and  $n$  unless some  $f_k$  is zero. This means that if  $\|\xi\| = 1$  we have an estimate:

$$\int_0^\infty \prod_{k=1}^n (F_k(e^{t\xi k}))^s \exp(-sta(-\xi)) dt \leq C \int_K \prod_{k=1}^n (F_k(e^{\xi k}))^s \exp(-sa(-\xi)) d\xi.$$

where  $K$  is a cylinder of radius one with axis  $\{t\xi : t \geq 0\}$ . We can then deduce (8) from (7) since  $f_k^* \leq F_k$ .

Finally (iv) implies (iii) (which implies (ii)) by just using polar coordinates. □

We can extend this result somewhat to certain multilinear analogues of maximal operators. Denote by  $L_0^+(0, \infty)$  the set of all nonnegative measurable functions on  $(0, \infty)$ . Let us say that a positively homogeneous (of degree 1) map  $T : \mathcal{E}_+^n \rightarrow L_0^+(0, \infty)$  is *n-quasi-sublinear* with constant  $C$  if for any  $k$  we have

$$(11) \quad T(f_1, \dots, f_{k-1}, (f_k + f'_k), f_{k+1}, \dots, f_n) \leq C(T(f_1, \dots, f_{k-1}, f_k, f_{k+1}, \dots, f_n) + T(f_1, \dots, f_{k-1}, f'_k, f_{k+1}, \dots, f_n)),$$

for all  $f_j, f'_j$ . Suppose  $T$  is *n-quasi-sublinear*. Then if we choose  $r$  so that  $2^{1/r-1} = C$  we can use the proof of the Aoki-Rolewicz theorem ([9],[14]) to deduce the existence of a constant  $C'$  so that for any  $1 \leq k \leq n$  and all  $m$  positive integers we have

$$T \left( f_1, \dots, f_{k-1}, \sum_{j=1}^m g_j, f_{k+1}, \dots, f_n \right) \leq C' \left( \sum_{j=1}^m T(f_1, \dots, f_{k-1}, g_j, f_{k+1}, \dots, f_n)^r \right)^{1/r},$$

for all  $f_j$  and  $g_j$ . Based on this it is easy to show the following, by exactly the same argument as in Theorem 3.2:

**Corollary 3.3.** *Suppose  $T : \mathcal{E}^n \rightarrow L_s(0, \infty)$  is n-quasi-sublinear with constant  $C = 2^{1/r-1}$  where  $0 < s \leq r$ , and that  $T$  is locally continuous. Let  $\Theta$  be a finite subset of  $\mathbf{R}_+^n$  and assume that there is a constant  $M$  so that*

$$\|T(\chi_{E_1}, \dots, \chi_{E_n})\|_{L_s} \leq M \inf_{\theta \in \Theta} \prod_{k=1}^n |E_k|^{\theta_k},$$

for all  $E_k$  of finite measure. If  $\mathbf{X}$  is an  $n$ -tuple of r.i. spaces satisfying (6), then there is a constant  $C$  so that for  $f_1, \dots, f_n \in \mathcal{E}$  we have

$$\|T(f_1, \dots, f_n)\|_{L_s} \leq CM \prod_{k=1}^n \|f_k\|_{X_k}.$$

We now consider versions of the Boyd interpolation theorem in this setting.

Consider the convex hull  $\text{co } \Theta$ ; We define the open convex hull  $\text{co}_0 \Theta$  to be the set of all  $\sum_{\theta \in \Theta} \alpha_\theta \theta$  where  $0 < \alpha_\theta < 1$  and  $\sum_{\theta \in \Theta} \alpha_\theta = 1$ . Then  $\text{co}_0 \Theta$  is the interior of  $\text{co } \Theta$  relative to the affine hyperplane it generates. We also define the *Boyd cube*  $B_{\mathbf{X}}$  of  $\mathbf{X}$  to be the set  $\prod_{k=1}^n [1/q_{X_k}, 1/p_{X_k}]$ , where  $p_{X_k}, q_{X_k}$  are the Boyd indices of  $X_k$ .

It will be convenient to introduce the following sublinear functional associated with  $B_{\mathbf{X}}$

$$(12) \quad b(\xi) = b_{\mathbf{X}}(\xi) = \max_{\phi \in B_{\mathbf{X}}} \langle \xi, \phi \rangle.$$

Let us first note a simple consequence of Theorem 3.2.

**Corollary 3.4.** *Suppose that  $\Theta$  is a finite subset of  $(\bar{\mathbf{R}}_+)^n$  and that  $\mathbf{X}$  satisfies the  $(\Theta, s)$ -interpolation condition. Then  $B_{\mathbf{X}} \cap \text{co } \Theta$  is nonempty.*

*Proof.* Suppose  $B_{\mathbf{X}}, \text{co } \Theta$  do not intersect. Then we can find  $\eta \in \mathbf{R}^n$  so that

$$\max_{\theta \in \Theta} \langle \eta, \theta \rangle < \min_{\phi \in B_{\mathbf{X}}} \langle \eta, \phi \rangle.$$

Thus  $a(\eta) = -b(-\eta) - 2\delta$  where  $\delta > 0$ . Now we refer to (1) to obtain for  $f \in \mathcal{E}$ ,

$$\prod_{k=1}^n \|D_{e^{-t\eta_k}} f_k^*\|_{X_k} \leq C \exp(t\delta + tb(-\eta)) \prod_{k=1}^n \|f_k^*\|_{X_k}$$

for  $t \geq 0$ . It follows from (6) that

$$\begin{aligned} \left( \int_{\mathbf{R}^n} \prod_{k=1}^n (f_k^*(e^{\xi_k + t\eta_k}))^s \exp(-sa(-\xi)) d\xi \right)^{1/s} &\leq C \exp(t\delta + tb(-\eta)) \prod_{k=1}^n \|f_k\|_{X_k} \\ &= C \exp(-ta(\eta) - t\delta) \prod_{k=1}^n \|f_k\|_{X_k}. \end{aligned}$$

Now  $a(-\xi) + ta(\eta) \geq a(-\xi + t\eta)$ . Thus we can reorganize to obtain

$$\left( \int_{\mathbf{R}^n} \prod_{k=1}^n (f_k^*(e^{\xi_k}))^s \exp(-sa(-\xi)) d\xi \right)^{1/s} \leq C \exp(-t\delta) \prod_{k=1}^n \|f_k\|_{X_k},$$

for every  $t \geq 0$  which is absurd. □

**Theorem 3.5.** *Suppose  $\Theta$  is a finite subset of  $(\overline{\mathbf{R}}_+)^n$ . Suppose  $\mathbf{X}$  is an  $n$ -tuple of r.i. spaces such that  $B_{\mathbf{X}} \cap \text{co } \Theta$  is a nonempty subset of  $\text{co}_0 \Theta$ . Then  $\mathbf{X}$  satisfies the  $(\Theta, s)$  interpolation condition provided there is a constant  $C$  so that if  $f_1, \dots, f_n \in \mathcal{E}$ ,*

$$(13) \quad \left( \int_H \prod_{k=1}^n (f_k^*(e^{\xi_k}))^s \exp(-sa(-\xi)) d\xi \right)^{1/s} \leq C \prod_{k=1}^n \|f_k\|_{X_k},$$

where  $H$  is the subspace of  $\mathbf{R}^n$  of all  $\xi$  such that  $\langle \xi, \theta \rangle$  is constant for all  $\theta \in \Theta$ .

If  $\Theta \subset \mathbf{R}_+^n$  and  $s\theta_k < 1$  for every  $\theta \in \Theta$  and  $1 \leq k \leq n$ , then inequality (13) is also necessary for  $\mathbf{X}$  to satisfy the  $(\Theta, s)$ -interpolation condition.

*Proof.* Let  $B_\epsilon = \{\xi : d(\xi, B_{\mathbf{X}}) \leq \epsilon\}$ . Our assumption on  $B_{\mathbf{X}}$  and a compactness argument give the existence of  $\epsilon > 0$  so that  $B_{2\epsilon} \cap P \subset \text{co } \Theta$ , where  $P$  is the affine plane generated by  $\Theta$ . We note that:

$$(14) \quad \max_{\phi \in B_{2\epsilon} \cap P} \langle \eta, \phi \rangle = \inf_{\xi \in H} a(\xi) + b(\eta - \xi) + 2\epsilon \|\eta - \xi\|.$$

To see this observe that the right hand side obviously dominates the left-hand side and is a sublinear functional. It is easy to check that if  $\langle \eta, \phi \rangle$  is dominated by the right-hand side then we have  $\phi \in B_{2\epsilon} \cap P$ .

We will also need (1) which implies that if  $f_1, \dots, f_n \in \mathcal{E}$ , then

$$\prod_{k=1}^n \|D_{e^{\eta_k}} f_k\|_{X_k} \leq C \exp(b(\eta) + \epsilon \|\eta\|) \prod_{k=1}^n \|f_k\|_{X_k}$$

for some constant  $C$ .

Now suppose  $\eta \in H^\perp$ . Then for a fixed  $\zeta \in H$  and  $f_1, \dots, f_n \in \mathcal{E}$  we have

$$\begin{aligned} & \left( \int_H \prod_{k=1}^n (f_k^*(e^{\xi_k + \eta_k}))^s \exp(-sa(-\xi - \eta)) d\xi \right)^{1/s} \\ &= \left( \int_H \prod_{k=1}^n (f_k^*(e^{\xi_k + \eta_k + \zeta_k}))^s \exp(-sa(-\xi - \zeta) - sa(-\eta)) d\xi \right)^{1/s} \\ &\leq \exp(-a(-\eta) + a(\zeta)) \left( \int_H \prod_{k=1}^n (f_k^*(e^{\xi_k + \eta_k + \zeta_k}))^s \exp(-sa(-\xi)) d\xi \right)^{1/s} \\ &\leq \exp(-a(-\eta) + a(\zeta)) \prod_{k=1}^n \|D_{e^{-\eta_k - \zeta_k}} f_k\|_{X_k} \\ &\leq C \exp(-a(-\eta) + a(\zeta) + b(-\eta - \zeta) + \epsilon \|\eta + \zeta\|) \prod_{k=1}^n \|f_k\|_{X_k}. \end{aligned}$$

At this point we use (14) and the fact that  $B_{2\epsilon} \cap P \subset \text{co } \Theta$  to show that

$$\inf_{\zeta \in H} (a(\zeta) + b(-\eta - \zeta) + \epsilon \|\eta + \zeta\|) \leq a(-\eta) - \epsilon \|\eta\|.$$

Thus we conclude that

$$\left( \int_H \prod_{k=1}^n (f_k^*(e^{\xi_k + \eta_k}))^s \exp(-sa(-\xi - \eta)) d\xi \right)^{1/s} \leq C \exp(-\epsilon \|\eta\|) \prod_{k=1}^n \|f_k\|_{X_k}.$$

Raise to the  $s^{\text{th}}$  power and integrate over  $\eta \in H^\perp$  to obtain (6). Hence  $\mathbf{X}$  satisfies the  $(\Theta, s)$ -interpolation condition.

The last statement follows from (8). □

We now specialize to the case when  $\Theta$  is a relatively large subset of  $(\overline{\mathbf{R}}_+)^n$ . Let us define the dimension of  $\Theta$ , denoted  $\dim \Theta$ , to be the dimension of the affine plane passing through all the points in  $\Theta$ . We say that  $\Theta$  is affinely independent if the conditions

$$\sum_{\theta \in \Theta} \lambda_\theta \theta = \mathbf{0} \quad \text{and} \quad \sum_{\theta \in \Theta} \lambda_\theta = 0$$

imply  $\lambda_\theta = 0, \forall \theta \in \Theta$ . Obviously if  $\Theta$  is affinely independent we have  $|\Theta| = 1 + \dim \Theta$ .

**Theorem 3.6.** *Suppose  $\dim \Theta = n$  (e.g. if  $\Theta$  is an affinely independent subset of  $(\overline{\mathbf{R}}_+)^n$  and  $|\Theta| = n + 1$ .) Suppose  $\mathbf{X}$  is an  $n$ -tuple of r.i. spaces such that  $B_{\mathbf{X}} \cap \text{co } \Theta$  is a nonempty subset of  $\text{co}_0 \Theta$ . Then  $\mathbf{X}$  satisfies the  $(\Theta, s)$  interpolation condition.*

*Proof.* In this case Theorem 3.5 applies with  $H = \{0\}$ . □

A more important case is the following:

**Theorem 3.7.** *Suppose that  $\dim \Theta = n - 1$  and that  $\Theta$  spans  $\mathbf{R}^n$ . Let  $0 < s \leq 1$  and suppose that  $\mathbf{X}$  is an  $n$ -tuple of r.i. spaces such that  $B_{\mathbf{X}} \cap \text{co } \Theta$  is a nonempty subset of  $\text{co}_0 \Theta$ . Pick a unique  $\sigma = (\sigma_k)_{k=1}^n$  so that  $\langle \sigma, \theta \rangle = 1$  for all  $\theta \in \Theta$ . Consider the following statements:*

- (i)  $\mathbf{X}$  satisfies the  $(\Theta, s)$  interpolation condition.
- (ii) There is a constant  $C$  such that if  $f_1, \dots, f_n \in \mathcal{E}$  we have

$$(15) \quad \left( \int_0^\infty x^{s-1} \prod_{k=1}^n (f_k^*(x^{\sigma_k}))^s dx \right)^{1/s} \leq C \prod_{k=1}^n \|f_k\|_{X_k}.$$

Then (ii) implies (i). Moreover, if  $\Theta \subset \mathbf{R}_+^n$  and  $s\theta_k < 1$  for every  $\theta \in \Theta$  and  $1 \leq k \leq n$ , then (i) and (ii) are equivalent.

Furthermore if  $\Theta \subset \mathbf{R}_+^n$  and if  $s(\sum_{k=1}^n \theta_k) \leq 1$  for every  $\theta \in \Theta$ , (i) and (ii) are also equivalent to:

(iii) There is a constant  $C$  so that if  $f_1, \dots, f_n \in \mathcal{E}$ ,

$$(16) \quad \left( \int_0^\infty x^{s-1} \prod_{\sigma_k \neq 0} |f_k(x^{\sigma_k})|^s dx \right)^{1/s} \leq C \prod_{\sigma_k \neq 0} \|f_k\|_{X_k}.$$

*Remarks.* The existence of  $\sigma$  follows from the fact that the plane generated by  $\Theta$  does not contain the origin. Note that the indices  $k$  for which  $\sigma_k = 0$  become *redundant* in the sense that that (15) can be rewritten as

$$\left( \int_0^\infty x^{s-1} \prod_{\sigma_k \neq 0} (f_k^*(x^{\sigma_k}))^s dx \right)^{1/s} \leq C \prod_{\sigma_k \neq 0} \|f_k\|_{X_k}.$$

Before we prove Theorem 3.7, let us illustrate the hypothesis on the Boyd indices, by considering the special but rather typical case when  $\text{co } \Theta$  is the intersection of a cube  $\prod_{k=1}^n [\alpha_k, \beta_k]$  with the plane  $\sum_{k=1}^n \theta_k = r^{-1}$ . In this case  $\sigma_k = r$  for all  $k$ . It may then be easily seen that the hypotheses on the Boyd indices are satisfied if we have both

$$(17) \quad \sum_{k=1}^n \frac{1}{q_{X_k}} \leq \frac{1}{r} \leq \sum_{k=1}^n \frac{1}{p_{X_k}}$$

and

$$(18) \quad \alpha_k < \frac{1}{q_{X_k}} \leq \frac{1}{p_{X_k}} < \beta_k$$

for all  $1 \leq k \leq n$ . However if for some  $l$  we have

$$(19) \quad \alpha_l + \sum_{k \neq l} \frac{1}{q_{X_k}} > \frac{1}{r}$$

then the lower bound condition on  $q_{X_l}^{-1}$  in (18) can be removed. Similarly if

$$(20) \quad \beta_l + \sum_{k \neq l} \frac{1}{p_{X_k}} < \frac{1}{r}$$

then the upper bound condition on  $p_{X_l}^{-1}$  can be removed.

*Proof.* The fact that (ii) implies (i) is an application of Theorem 3.5. Indeed, in this case  $H$  is one-dimensional, say  $H = \{t\sigma\}_{t \in \mathbf{R}}$ . Then equation (13) becomes

$$\left( \int_{-\infty}^{+\infty} \prod_{k=1}^n (f_k^*(e^{t\sigma_k}))^s e^{-st} dt \right)^{1/s} \leq C \prod_{k=1}^n \|f_k\|_{X_k}$$

which reduces to (15) by substituting  $x = e^t$ . The converse statement follows from Theorem 3.5.

We now prove that (i) implies (iii) under the extra hypothesis  $s(\sum_{k=1}^n \theta_k) \leq 1$  for every  $\theta \in \Theta$ . Define a map  $T : \mathcal{E}^n \rightarrow L_s((0, \infty) \times (0, 1))$  by setting

$$T(f_1, \dots, f_n)(x, y) = x^{1-1/s} \prod_{\sigma_k \neq 0} f_k(x^{\sigma_k}) \prod_{\sigma_l = 0} f_l(y).$$

We will show that  $T$  is  $\Theta$ -admissible.

Suppose  $(E_k)$  are sets of finite measure. Let  $F = \{x : x^{\sigma_k} \in E, \forall \sigma_k \neq 0\}$  and let  $G = [0, 1] \cap \cap_{\sigma_k=0} E_k$ . Then we have

$$T(\chi_{E_1}, \dots, \chi_{E_n})(x, y) = x^{1-1/s} \chi_F(x) \chi_G(y)$$

and therefore

$$\|T(\chi_{E_1}, \dots, \chi_{E_n})\| = \left( \int_F x^{s-1} dx \right)^{1/s} |G|^{1/s}.$$

Now suppose  $\theta \in \Theta$ . Let  $r = (\sum_{\sigma_k \neq 0} \theta_k)^{-1}$ . Clearly  $s \leq r$ . We have by Lemma 3.1

$$\begin{aligned} \left( \int_F x^{s-1} dx \right)^{1/s} &\leq r^{1/r} s^{-1/s} \left( \int_F x^{r-1} dx \right)^{1/r} \\ &\leq r^{1/r} s^{-1/s} \prod_{j \in J} \left( \int_F x^{\sigma_k-1} dx \right)^{\theta_k} \\ &\leq r^{1/r} s^{-1/s} \prod_{k \in J} |\sigma_k|^{-\theta_k} |E_k|^{\theta_k}. \end{aligned}$$

On the other hand since  $|G| \leq 1$  and  $\sum_{\sigma_k \neq 0} \theta_k \leq s^{-1}$ ,

$$|G|^{1/s} \leq \prod_{\sigma_k \neq 0} |E_k|^{\theta_k}.$$

Thus  $T$  is  $\Theta$ -admissible and hence  $T$  extends to a bounded  $n$ -linear form on  $X_1 \times \dots \times X_n$ . Letting  $f_k = \chi_{[0,1]}$  if  $\sigma_k = 0$  and restricting gives (16).

Now it is clear that (iii) implies (ii) and so the proof is complete. □

**Corollary 3.8.** *Suppose under the hypotheses of Theorem 3.7 we also have that for some fixed  $r$*

$$\sum_{k=1}^n \theta_k = \frac{1}{r}$$

for every  $\theta \in \Theta$ . Then  $\mathbf{X}$  has the  $(\Theta, s)$  interpolation condition if

$$(21) \quad \left( \int_0^\infty x^{s/r-1} \prod_{k=1}^n (f_k^*(x))^s dx \right)^{1/s} \leq C \prod_{k=1}^n \|f_k\|_{X_k}.$$

In particular if  $r = s$  then  $\mathbf{X}$  has the  $(\Theta, s)$  interpolation condition if and only if  $X_1 \cdots X_n \subset L_s$  where  $X_1 \cdots X_n = \{f_1 \dots f_n; f_k \in X_k\}$ .

*Proof.* In this case  $\sigma_k = r$  for all  $k$  and (21) is obtained by a simple change of variables from (15). □

Finally let us note an unusual case which can arise:

**Theorem 3.9.** *Suppose  $\Theta \subset \mathbf{R}_+^n$ ,  $\dim \Theta = n - 1$  and  $\Theta$  does not span  $\mathbf{R}^n$ . Suppose  $\mathbf{X}$  is an  $n$ -tuple of r.i. spaces such that  $B_{\mathbf{X}} \cap \text{co } \Theta$  is a nonempty subset of  $\text{co}_0 \Theta$ . Let  $\sigma = (\sigma_k)_{k=1}^n$  be chosen so that  $\langle \sigma, \theta \rangle = 0$  for all  $\theta \in \Theta$ . Assume  $s > 0$  is such that  $s\theta_k < 1$  for every  $\theta \in \Theta$  and  $1 \leq k \leq n$ . Then the following are equivalent:*

- (i)  $\mathbf{X}$  satisfies the  $(\Theta, s)$  interpolation condition.
- (ii) There is a constant  $C$  so that for  $f_1, \dots, f_n \in \mathcal{E}$  we have

$$(22) \quad \left( \int_0^\infty x^{-1} \prod_{k=1}^n (f_k^*(x^{\sigma_k}))^s dx \right)^{1/s} \leq C \prod_{k=1}^n \|f_k\|_{X_k}.$$

We omit the proof which is similar to the that of Theorem 3.7.

#### 4. The inhomogeneous multilinear Boyd theorem and applications

Suppose that  $\Theta$  is a finite subset of  $(\overline{\mathbf{R}}_+)^n$  and that  $\theta \rightarrow r_\theta$  is a map from  $\Theta$  to  $\mathbf{R}_+$ . We denote  $\phi_\theta = (\theta, r_\theta)$  and  $\Phi = \{\phi_\theta : \theta \in \Theta\}$ . Clearly  $\Phi \subset \mathbf{R}^{n+1}$ . Now consider the case when we are given a map  $T : \mathcal{E}^n \rightarrow L_0(0, \infty)$ , which we assume to be locally continuous. We will say that  $T$  satisfies a weak-type  $(\theta, r_\theta)$  estimate if there exists  $M > 0$  so that if  $E_1, \dots, E_n$  are sets of finite measure then

$$(23) \quad \|T(\chi_{E_1}, \dots, \chi_{E_n})\|_{L_{r,\infty}} \leq M \prod_{k=1}^n |E_k|^{\theta_k}.$$

We now give a version of the Boyd interpolation theorem for this setting which follows almost immediately from Theorem 3.5. For simplicity we shall only treat the most important case.

**Theorem 4.1.** *Suppose  $\Theta$  is a subset of  $(\overline{\mathbf{R}}_+)^n$  with  $|\Theta| = n + 1$  and  $\dim \Theta = n$ . Suppose for each  $\theta$  we have  $0 < r_\theta \leq \infty$ . Let  $\sigma \in \mathbf{R}^n$  be the unique solution of the equation*

$$\langle \sigma, \theta \rangle = \frac{1}{r_\theta} + \tau$$

where  $\tau$  is independent of  $\theta$ . Let  $\mathbf{X}$  be an  $n$ -tuple of r.i. spaces and suppose  $Y$  is a maximal r.i. space which is  $s$ -convex for some  $s > 0$ . Suppose the Boyd cube  $B_{\mathbf{X}} \times [1/q_Y, 1/p_Y]$  intersects  $\text{co } \Phi$  in a non-empty subset of  $\text{co}_0 \Phi$ .

Then in order that every locally continuous  $n$ -linear  $T : \mathcal{E}^n \rightarrow L_0(0, \infty)$ , which satisfies the weak type  $(\theta, r_\theta)$  estimate (23) for  $\theta \in \Theta$ , extends to a bounded  $n$ -linear map  $T : \prod_{k=1}^n X_k \rightarrow Y$  (with norm a multiple of  $M$ ), it is sufficient that there is a constant  $C$  so that if  $f_1, \dots, f_n \in \mathcal{E}$  then

$$(24) \quad \|x^\tau \prod_{k=1}^n f_k^*(x^{\sigma_k})\|_Y \leq C \prod_{k=1}^n \|f_k\|_{X_k}.$$

If in addition we have

$$0 < \frac{1}{r_\theta} \leq \sum_{k=1}^n \theta_k$$

for every  $\theta \in \Theta$ , then (24) is also necessary and is equivalent to the condition that there exists a constant  $C$  so that for  $f_1, \dots, f_n \in \mathcal{E}$  we have

$$(25) \quad \|x^\tau \prod_{\sigma_k \neq 0} f_k(x^{\sigma_k})\|_Y \leq C \prod_{\sigma_k \neq 0} \|f_k\|_{X_k}.$$

*Remark.* The same conclusions can be obtained if  $|\Theta| > n + 1$  provided there is a solution of the equation  $\langle \sigma, \theta \rangle = r_\theta^{-1} + \tau$ .

*Proof.* We first choose  $s > 0$  small enough so that  $Y$  is  $s$ -convex,  $s < r_\theta$  for all  $\theta \in \Theta$ , and  $\tau + \frac{1}{s} > 0$ . Next define  $X_{n+1}$  to be the space of all  $f \in L_0(0, \infty)$  so that  $fg \in L_s(0, \infty)$  for all  $g \in Y$  with the quasi-norm

$$\|f\|_{X_{n+1}} = \sup_{\|g\|_Y \leq 1} \|fg\|_{L_s}.$$

Since  $Y$  is both maximal and  $s$ -convex we obtain that  $g \in Y$  if and only if  $\sup\{\|fg\|_{L_s} : \|f\|_{X_{n+1}} \leq 1\}$  is finite and furthermore there is a constant  $C$  so that  $\|f\|_Y \leq \sup\{\|fg\|_{L_s} : \|f\|_{X_{n+1}} \leq 1\}$ . (If  $Y$  is  $s$ -convex with constant one then  $C = 1$ ; this is easily seen by noting that  $Y^s$  is a Banach r.i. space and  $X_{n+1}^s$  is simply the Köthe dual space; in general we can always renorm  $Y$  to have  $s$ -convexity constant one.) It easy to calculate the Boyd indices of  $X_{n+1}$ ; these are given by

$$\frac{1}{p_{X_{n+1}}} = \frac{1}{s} - \frac{1}{q_Y}, \quad \frac{1}{q_{X_{n+1}}} = \frac{1}{s} - \frac{1}{p_Y}.$$



We refer to [12] for similar calculations for dual spaces.

Now if  $T$  satisfies the weak-type  $(\theta, r_\theta)$  estimate (23) for every  $\theta \in \Theta$ , then we consider the map  $T' : \mathcal{E}^{n+1} \rightarrow L_s(0, \infty)$  defined by

$$T'(f_1, \dots, f_{n+1}) = T(f_1, \dots, f_n) f_{n+1}.$$

If  $E_1, \dots, E_{n+1}$  are sets of finite measure then

$$\begin{aligned} \|T'(\chi_{E_1}, \dots, \chi_{E_{n+1}})\|_{L_s} &\leq \left( \int_{E_{n+1}} |T(\chi_{E_1}, \dots, \chi_{E_n})|^s dx \right)^{1/s} \\ &\leq M \left( \frac{r_\theta}{r_\theta - s} \right)^{1/s} |E_{n+1}|^{1/s-1/r_\theta} \prod_{k=1}^n |E_k|^{\theta_k} \end{aligned}$$

for every  $\theta \in \Theta$ . Thus if we let  $\psi_\theta = (\theta, \frac{1}{s} - \frac{1}{r_\theta})$  and  $\Psi = \{\psi_\theta : \theta \in \Theta\}$  then  $T'$  is  $(\Psi, s)$ -admissible. It is clear from our discussion of  $X_{n+1}$  that we only need to show that  $T'$  extends to a bounded  $n$ -linear map on  $X_1 \times \dots \times X_{n+1}$ .

We now use Theorem 3.7. We first argue that  $\Psi$  is linearly independent in  $\mathbf{R}^{n+1}$ . Indeed from the definition of  $\tau$  this is equivalent to the linear independence of the points  $(\theta, \tau + \frac{1}{s})$  which follows from the affine independence of  $\Theta$ . We will show that the  $(n+1)$ -tuple  $(X_1, \dots, X_{n+1})$  has the  $(\Psi, s)$ -interpolation condition. We note that our hypotheses on the Boyd indices of  $\mathbf{X}$  and  $Y$  imply that the hypotheses on the Boyd indices for Theorem 3.6 hold. Define  $\sigma'_k = \sigma_k(\tau + \frac{1}{s})^{-1}$  for  $k \leq n$  and  $\sigma'_{n+1} = (\tau + \frac{1}{s})^{-1}$ . Then

$$\langle \sigma', \psi_\theta \rangle = (\tau + \frac{1}{s})^{-1} (\tau + \frac{1}{r_\theta}) + (\tau + \frac{1}{s})^{-1} (\frac{1}{s} - \frac{1}{r_\theta}) = 1.$$

Now if  $f_1, \dots, f_{n+1} \in \mathcal{E}^{n+1}$ , we have

$$\begin{aligned} \left( \int_0^\infty x^{s-1} \prod_{k=1}^{n+1} |f_k^*(x^{\sigma'_k})|^s dx \right)^{1/s} &= \\ \left( \tau + \frac{1}{s} \right)^{1/s} \left( \int_0^\infty x^{s\tau} |f_{n+1}^*(x)|^s \prod_{k=1}^n |f_k^*(x^{\sigma_k})|^s dx \right)^{1/s}. \end{aligned}$$

Now it is clear that if we assume (24) then we obtain (15) in Theorem 3.7 and so  $(X_1, \dots, X_{n+1})$  satisfies the interpolation condition  $(\Theta, s)$ .

For the second part we construct the map  $T : \mathcal{E}^n \rightarrow L_0((0, \infty) \times (0, 1))$  by

$$T(f_1, \dots, f_n)(x, y) = x^\tau \prod_{\sigma_k \neq 0} f_k(x^{\sigma_k}) \prod_{\sigma_j = 0} f_j(y).$$

Let  $u_\theta = (\sum_{\sigma_k \neq 0} \theta_k)^{-1}$  so that  $r_\theta \leq u_\theta$ . Then by arguments similar to those for Theorem 3.7 we have that if  $E_1, \dots, E_n$  are measurable sets of finite measure,

$$\|T(\chi_{E_1}, \dots, \chi_{E_n})\|_{L_{r_\theta}} \leq r_\theta^{-1/r_\theta} u_\theta^{1/u_\theta} \prod_{k=1}^n |E_k|^{\theta_k}.$$

Our hypotheses then guarantee that  $T$  maps  $X_1 \times \dots \times X_n$  into  $Y$  i.e. we have (25) and hence also (24). □

Let us isolate a simple special case:

**Corollary 4.2.** *Suppose that in the preceding theorem we have*

$$\sum_{k=1}^n \theta_k = \frac{1}{r_\theta}$$

for every  $\theta \in \Theta$ . Then (25) is equivalent to the inclusion  $X_1 \cdots X_n \subset Y$ , where  $X_1 \cdots X_n$  is the set of all products  $f_1 \cdots f_n$  with  $f_k \in X_k$ .

*Proof.* We need only to observe that in this case  $\sigma_k = 1$  for every  $k$  and  $\tau = 0$ . □

We next point out that under certain hypotheses, we can replace (24) with an alternative criterion:

**Corollary 4.3.** *Suppose that in Theorem 4.1,  $\tilde{Y}$  is a carrier space for  $Y$  with the property that  $\|D_a\|_{\tilde{Y}} \leq C_0 a^\rho$  for all  $0 < a < 1$  where  $\rho > 0$  and that  $\tilde{Y}$  is  $s$ -convex for some  $s > 0$ . Then the sufficient condition (24) can be replaced by:*

$$(26) \quad \|x^\tau \prod_{k=1}^n f_k^*(x^{\sigma_k})\|_{\tilde{Y}} \leq C \prod_{k=1}^n \|f_k\|_{X_k},$$

for  $f_1, \dots, f_n \in \mathcal{E}$ .

*Proof.* We note that in the proof of Theorem 24 we can take  $s$  small enough so  $\tilde{Y}$  is  $s$ -convex. Suppose  $f_1, \dots, f_n \in \mathcal{E}$ . Let  $\varphi(x) = x^\tau \prod_{k=1}^n f_k^*(x^{\sigma_k})$ . By assumption  $\|\varphi\|_{\tilde{Y}} \leq C \prod_{k=1}^n \|f_k\|_{X_k}$ . Now let  $\psi$  be defined by

$$\psi(x) = \left( \int_x^\infty \varphi(y)^s \frac{dy}{y} \right)^{1/s}.$$

By the  $s$ -convexity of  $\tilde{Y}$  we obtain that

$$\|\psi\|_{\tilde{Y}} \leq M \left( \int_0^1 a^{s\rho-1} da \right)^{1/s} \|\varphi\|_{\tilde{Y}}$$

so that we have an estimate

$$\|\psi\|_{\tilde{Y}} \leq C_1 \|\varphi\|_{\tilde{Y}}.$$

However  $\psi$  is decreasing and so  $\|\psi\|_Y \leq C_1 \|\varphi\|_{\tilde{Y}}$ . Now if  $f_{n+1} \in \mathcal{E}$  we have that

$$\left( \int_0^\infty (f_{n+1}^*(x))^s \varphi(x)^s dx \right)^{1/s} \leq C C_1 \prod_{k=1}^{n+1} \|f_k\|_{X_k}$$

and the proof is completed in the same way. □

At this point we note that we can use Corollary 3.3 to extend this result to  $n$ -quasi-sublinear maps.

**Corollary 4.4.** *Assume that  $X_1, \dots, X_n, Y$  satisfy (24). Suppose  $T : \mathcal{E}^n \rightarrow L_0(0, \infty)$  is  $n$ -quasi-sublinear, locally continuous, and satisfies the weak-type  $(\theta, r_\theta)$ -inequality (23) for every  $\theta \in \Theta$ . Then we have the estimate*

$$\|T(f_1, \dots, f_n)\|_Y \leq CM \prod_{k=1}^n \|f_k\|_{X_k}$$

for  $f_1, \dots, f_n \in \mathcal{E}$ .

We omit the details of the proof. The key point to note is that we should choose  $s$  in the argument for Theorem 4.1 above sufficiently small so that  $2^{1/s-1} \geq C$  where  $C$  is the constant in (11).

It is also worth noting that we can give a similar result to Theorem 4.1 in the case when  $\Theta$  fails to be affinely independent. This case is somewhat degenerate. For example in the case  $n = 1$  it applies to linear operators which satisfy weak type estimates  $(p, q_1)$  and  $(p, q_2)$  where  $q_1 \neq q_2$ .

**Theorem 4.5.** *Suppose  $\Theta$  is an affinely dependent subset of  $(\overline{\mathbf{R}}_+)^n$  with  $|\Theta| = n + 1$ . Suppose for each  $\theta$  we have  $0 < r_\theta \leq \infty$  and that the set  $\Phi = \{(\theta, r_\theta^{-1}) : \theta \in \Theta\}$  is linearly independent in  $\mathbf{R}^{n+1}$ . Choose  $\sigma \in \mathbf{R}^n$  so that  $\langle \sigma, \theta \rangle = 1$  for all  $\theta \in \Theta$ . Let  $\mathbf{X}$  be an  $n$ -tuple of r.i. spaces and suppose  $Y$  is a maximal r.i. space. Let  $r = \min_{\theta \in \Theta} r_\theta$  and suppose  $0 < s \leq 1$  is such that  $s < 1$  if  $r = 1$  and  $s \leq r$  otherwise. Suppose also the Boyd cube  $B_{\mathbf{X}} \times [1/q_Y, 1/p_Y]$  intersects  $\text{co } \Phi$  in a non-empty subset of  $\text{co } \Phi$ .*

*Then, in order that every locally continuous  $n$ -linear  $T : \mathcal{E}^n \rightarrow L_0(0, \infty)$ , which satisfies the weak type  $(\theta, r_\theta)$  estimate (23) for  $\theta \in \Theta$ , extends to a bounded  $n$ -linear map  $T : \prod_{k=1}^n X_k \rightarrow Y$  (with norm a multiple of  $M$ ), it is sufficient that there exists a constant  $C$  so that*

$$(27) \quad \left( \int_0^\infty x^{s-1} \prod_{k=1}^n (f_k^*(x^{\sigma_k}))^s dx \right)^{1/s} \leq C \prod_{k=1}^n \|f_k\|_{X_k},$$

for  $f_1, \dots, f_n \in \mathcal{E}$ .

*Remark.* The existence and uniqueness of  $\sigma$  is a consequence of our hypotheses, since  $\Theta$  generates a plane of dimension  $n - 1$  which cannot be a linear subspace.

*Proof.* Our hypotheses are such that the space  $L_{r,\infty}$  is  $s$ -normable. In this case the convex set  $\Phi$  generates a plane containing the line in the direction parallel to the basis vector  $e_{n+1}$ .

We first prove the result when  $Y = L_{t,\infty}$  for some  $t$ . By the above remark we have  $t > r$ . Let  $X_{n+1} = L_{u,\infty}$  where  $\frac{1}{t} + \frac{1}{u} = \frac{1}{r}$ . Let  $\psi_\theta = (\theta, \frac{1}{r} - \frac{1}{r_\theta}) \in \mathbf{R}^{n+1}$  and  $\Psi = \{\psi_\theta : \theta \in \Theta\}$ . Now it is clear that (27) implies that the  $(n + 1)$ -tuple  $(X_1, \dots, X_{n+1})$  satisfies the conditions of Theorem 3.7 for the  $(\Psi, s)$ -interpolation condition. We apply this to the map  $T' : \mathcal{E}^{n+1} \rightarrow L_{r,\infty}$  where  $T'(f_1, \dots, f_{n+1}) = T(f_1, \dots, f_n) f_{n+1}$ . A routine calculation gives

$$\|T'(\chi_{E_1}, \dots, \chi_{E_{n+1}})\|_{L_{r,\infty}} \leq CM|E|^{1/r-1/r_\theta} \prod_{k=1}^n |E_k|^{\theta_k}.$$

Hence we have the estimate

$$\|T(f_1, \dots, f_n) f_{n+1}\|_{L_{r,\infty}} \leq CM \|f_{n+1}\|_{L_{u,\infty}} \prod_{k=1}^n \|f_k\|_{X_k}.$$

This implies the estimate

$$\|T(f_1, \dots, f_n)\|_{L_{t,\infty}} \leq CM \prod_{k=1}^n \|f_k\|_{X_k}$$

by simply considering  $f_{n+1} = \chi_E$  for  $E$  a set of finite measure. We have now proved our claim.

Next we consider the general case. By our assumptions on  $\Phi$  we may find  $t < p_Y \leq q_Y < u$  so that both  $(X_1, \dots, X_n, L_{t,\infty})$  and  $(X_1, \dots, X_n, L_{u,\infty})$  satisfy the interior condition on the Boyd indices. Hence  $T$  maps  $X_1 \times \dots \times X_n$  boundedly into  $L_{t,\infty} \cap L_{u,\infty}$  with norm a multiple of  $M$ . But it is easy to calculate from the Boyd indices that  $L_{t,\infty} \cap L_{u,\infty} \subset Y$ . □

The theorems below extend the classical Marcinkiewicz interpolation theorem to the multilinear setting.

**Theorem 4.6.** *Let  $0 < p_{jk} \leq \infty$  for  $1 \leq j \leq n + 1$  and  $1 \leq k \leq n$ , and also let  $0 < p_j \leq \infty$  for  $1 \leq j \leq n + 1$ . Suppose that a locally continuous  $n$ -linear map  $T : \mathcal{E}^n \rightarrow L_0$  satisfies*

$$\|T(\chi_{E_1}, \dots, \chi_{E_n})\|_{L_{p_j,\infty}} \leq M|E_1|^{1/p_{j1}} \dots |E_n|^{1/p_{jn}}$$

for all sets  $E_j$  of finite measure and all  $1 \leq j \leq n + 1$ . Assume that the system below

$$\begin{pmatrix} 1/p_{11} & 1/p_{12} & \dots & 1/p_{1n} & 1 \\ 1/p_{21} & 1/p_{22} & \dots & 1/p_{2n} & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1/p_{n1} & 1/p_{n2} & \dots & 1/p_{nn} & 1 \\ 1/p_{(n+1)1} & 1/p_{(n+1)2} & \dots & 1/p_{(n+1)n} & 1 \end{pmatrix} \begin{pmatrix} \sigma_1 \\ \sigma_2 \\ \vdots \\ \sigma_n \\ -\tau \end{pmatrix} = \begin{pmatrix} 1/p_1 \\ 1/p_2 \\ \vdots \\ 1/p_n \\ 1/p_{n+1} \end{pmatrix},$$

has a unique solution  $(\sigma_1, \dots, \sigma_n, -\tau) \in \mathbf{R}^{n+1}$  with not all  $\sigma_j = 0$ . Suppose that  $(1/q_1, \dots, 1/q_n, 1/q)$  lies in the open convex hull of the points  $(1/p_{j1}, \dots, 1/p_{jn}, 1/p_j)$  in  $\mathbf{R}^{n+1}$  and let  $0 < t_k, t \leq \infty$  satisfy

$$(28) \quad \sum_{\substack{1 \leq k \leq n \\ \sigma_k \neq 0}} \frac{1}{t_k} \geq \frac{1}{t}.$$

Then  $T$  extends to a bounded  $n$ -linear map  $T : \prod_{k=1}^n L_{q_k, t_k} \rightarrow L_{q, t}$  with norm a multiple of  $M$ .

*Remark.* We remark that the existence of the unique solution in the linear system of Theorem 4.6 is equivalent to the condition that the  $n + 1$  points  $\theta_j = (1/p_{jk})_{k=1}^n$  are affinely independent in  $\mathbf{R}^n$ . We also note that as in Corollary 4.4 the result above is valid for  $n$ -quasi-sublinear maps.

*Proof.* We clearly only need to consider the case of equality in (28). It is clear that the Boyd index assumption of Theorem 4.1 is satisfied. Clearly we have

$$\sum_{k=1}^n \frac{\sigma_k}{q_k} = \tau + \frac{1}{q}.$$

Hence if  $f_1, \dots, f_n \in \mathcal{E}$  we have

$$x^{1/q+\tau} \prod_{k=1}^n f_k^*(x^{\sigma_k}) = \prod_{k=1}^n x^{\sigma_k/q_k} f_k^*(x^{\sigma_k}).$$

Let  $F(x) = x^\tau \prod_{\sigma_k \neq 0} f_k^*(x^{\sigma_k})$ . Then

$$\left( \int_0^\infty (x^{1/q} F(x))^t \frac{dx}{x} \right)^{1/t} \leq C \prod_{\sigma_k \neq 0} \left( \int_0^\infty (x^{\sigma_k/q_k} f_k^*(x^{\sigma_k}))^{t_k} \frac{dx}{x} \right)^{1/t_k}.$$

Now if  $\sigma_k \neq 0$

$$\left( \int_0^\infty (x^{\sigma_k/q_k} f_k^*(x^{\sigma_k}))^{t_k} \frac{dx}{x} \right)^{1/t_k} = |\sigma_k|^{-1} \|f\|_{L_{q_k, t_k}}.$$

Thus we have an estimate

$$\left( \int_0^\infty (x^{1/q} F(x))^t \frac{dx}{x} \right)^{1/t} \leq C \prod_{\sigma_k \neq 0} \|f_k\|_{L_{q_k, t_k}}.$$

In view of Corollary 4.3 this completes the proof. □

There is a version of the above result for the degenerate case corresponding to Theorem 4.5:

**Theorem 4.7.** *Let  $0 < p_{jk} \leq \infty$  for  $1 \leq j \leq n + 1$  and  $1 \leq k \leq n$ , and let  $0 < p_j \leq \infty$  for  $1 \leq j \leq n + 1$ . Suppose that a locally continuous  $n$ -linear map  $T : \mathcal{E}^n \rightarrow L_0$  satisfies*

$$\|T(\chi_{E_1}, \dots, \chi_{E_n})\|_{L_{p_j, \infty}} \leq M |E_1|^{1/p_{j1}} \dots |E_n|^{1/p_{jn}}$$

for all subsets  $E_k$  of finite measure and all  $1 \leq j \leq n + 1$ . Assume that the  $n + 1$  points  $\theta_j = (1/p_{jk})_{k=1}^n$  are affinely dependent in  $\mathbf{R}^n$ , but the points  $(\theta_j, 1/p_j)$  are linearly independent in  $\mathbf{R}^{n+1}$ . Suppose that  $(1/q_1, \dots, 1/q_n, 1/q)$  lies in the open convex hull of the points  $(1/p_{j1}, \dots, 1/p_{jn}, 1/p_j)$  in  $\mathbf{R}^{n+1}$ . Let  $r = \min_{1 \leq j \leq n+1} p_j$  and  $0 < t_k, t \leq \infty$  satisfy

$$(29) \quad \sum_{\substack{1 \leq k \leq n \\ \sigma_k \neq 0}} \frac{1}{t_k} \begin{cases} > 1 & \text{if } r = 1, \\ \geq \frac{1}{r} & \text{if } r \neq 1, \end{cases}$$

where  $\{\sigma_k\}_{k=1}^n$  are the unique solutions of the system

$$\sum_{k=1}^n \frac{\sigma_k}{p_{jk}} = 1, \quad 1 \leq j \leq n + 1.$$

Then  $T$  extends to a bounded  $n$ -linear map  $T : \prod_{k=1}^n L_{q_k, t_k} \rightarrow L_{q, t}$  with norm a multiple of  $M$ .

*Proof.* This is deduced from Theorem 4.5. It is clear our hypotheses guarantee the appropriate conditions on the Boyd indices. Pick any  $0 < s \leq 1$  so that  $s \leq r$  if  $r \neq 1$  and  $s < 1$  otherwise with

$$\frac{1}{s} \geq \sum_{\sigma_k \neq 0} \frac{1}{t_k}.$$

It then suffices to verify (27) in Theorem 4.5. To do this we can clearly suppose that

$$\frac{1}{s} = \sum_{\sigma_k \neq 0} \frac{1}{t_k}.$$

Suppose  $f_1, \dots, f_n \in \mathcal{E}$  and set

$$F(x) = x \prod_{\sigma_k \neq 0} f_k^*(x^{\sigma_k}).$$

Then

$$F(x) = \prod_{\sigma_k \neq 0} x^{\sigma_k/q_k} f_k^*(x^{\sigma_k})$$

and so

$$\left( \int_0^\infty F(x)^s \frac{dx}{x} \right)^{1/s} \leq \prod_{\sigma_k \neq 0} |\sigma_k|^{-1} \|f_k\|_{L_{q_k, t_k}}.$$

This establishes (27) and completes the proof. □

### 5. Examples and applications

In this section we discuss some examples of multilinear interpolation. For simplicity we restrict ourselves to bilinear and trilinear examples.

*Example 5.1.* (Young’s inequality and O’Neil’s inequality) On a locally compact abelian group consider the bilinear operator  $(f, g) \rightarrow f * g$ , where  $*$  denotes convolution. Let  $H$  denote the closed triangle in  $\mathbf{R}^3$  with vertices  $(1, 0, 0)$ ,  $(0, 1, 0)$ , and  $(1, 1, 1)$ . The well known Young’s inequality says that

$$(30) \quad \|f * g\|_{L_r} \leq C \|f\|_{L_p} \|g\|_{L_q}$$

holds if the point  $(1/p, 1/q, 1/r)$  lies in the closure of the triangle  $H$ .

The three trivial estimates

$$\begin{aligned} \|f * g\|_{L_1} &\leq \|f\|_{L_1} \|g\|_{L_1}, \\ \|f * g\|_{L_\infty} &\leq \|f\|_{L_1} \|g\|_{L_\infty}, \text{ and} \\ \|f * g\|_{L_\infty} &\leq \|f\|_{L_\infty} \|g\|_{L_1} \end{aligned}$$

give (30) on the interior of  $H$ . The estimates on the sides follow from bilinear complex interpolation.

Applying Theorem 4.6 in the situation above we obtain O’Neil’s inequality. If the point  $(1/p, 1/q, 1/r)$  lies in the interior of the triangle  $H$  and  $0 < s_1, s_2 \leq \infty$  and  $1/s = 1/s_1 + 1/s_2$ , then

$$(31) \quad \|f * g\|_{L_{r,s}} \leq C \|f\|_{L_{p,s_1}} \|g\|_{L_{q,s_2}}.$$

The special case  $s_1 = p, s_2 = \infty$  is of particular interest. Observe that if  $(1/p, 1/q, 1/r)$  lies in the interior of  $H$ , then  $1/p + 1/q = 1/r + 1$ , from which it follows that  $p < r$ , which in turn implies that

$$\|f * g\|_{L_r} \leq C \|f * g\|_{L_{r,p}} \leq C \|f\|_{L_p} \|g\|_{L_{q,\infty}}.$$

The inequality above provides a sharpening of Young’s inequality since the space  $L_q$  is replaced by  $L^{q,\infty}$ .

More generally we can use Theorem 4.1 to obtain the following result:

**Theorem 5.2.** *Suppose  $X, Y, Z$  are r.i. spaces whose Boyd indices satisfy the conditions*

$$1 < p_X, p_Y, p_Z, q_X, q_Y, q_Z < \infty,$$

$$\frac{1}{p_X} + \frac{1}{p_Y} \geq 1 + \frac{1}{q_Z},$$

and

$$\frac{1}{q_X} + \frac{1}{q_Y} \leq 1 + \frac{1}{p_Z}.$$

Assume that  $Z$  is maximal and  $s$ -convex for some  $s > 0$ . Then  $(f, g) \rightarrow f * g$  maps  $X \times Y$  to  $Z$  provided the map  $(f, g) \rightarrow xf(x)g(x)$  maps  $X(0, \infty) \times Y(0, \infty)$  to  $Z(0, \infty)$ .

*Remark.* Of course we can state this theorem with less stringent requirements on the Boyd indices, namely that the Boyd cube intersects  $H$  in a subset of its relative interior. As in the discussion in the remarks after Theorem 3.7 this can be illustrated. We can allow for example  $p_X \leq 1$  provided  $q_Y < p_Z$ , and  $q_X = \infty$  is permissible provided  $p_Y^{-1} < 1 + q_Z^{-1}$ . Similarly  $p_Z \leq 1$  is permissible if  $p_X^{-1} + p_Y^{-1} < 2$  and  $q_Z = \infty$  is permissible if  $q_X^{-1} + q_Y^{-1} > 1$ .

*Example 5.3.* Fix three numbers  $0 < \alpha, \beta, \gamma < n$  such that  $\alpha + \beta > n, \beta + \gamma > n$  and  $\gamma + \alpha > n$ . Consider now the trilinear fractional integral form

$$I_{\alpha,\beta,\gamma}(f, g, h) = \int_{\mathbf{R}^n} \int_{\mathbf{R}^n} \int_{\mathbf{R}^n} f(x)g(y)h(z)|x - y|^{-\alpha}|y - z|^{-\beta}|z - x|^{-\gamma} dx dy dz.$$

We claim that the following inequality is valid

$$|I_{\alpha,\beta,\gamma}(f, g, h)| \leq C \|f\|_{L_p} \|g\|_{L_q} \|h\|_{L_r}$$

if and only if

$$(32) \quad \frac{1}{p} + \frac{1}{q} + \frac{1}{r} + \frac{\alpha + \beta + \gamma}{n} = 3,$$

$$1 < p, q, r < \infty, \quad \text{and} \quad \frac{1}{p} + \frac{1}{q} + \frac{1}{r} > 1.$$

Note that (32) requires  $\alpha + \beta + \gamma < 2n$ .

Examples can be given to prove the necessity of the conditions on the indices above. Let us prove here the sufficiency. The assumptions  $\alpha + \beta > n, \beta + \gamma > n$ , and  $\gamma + \alpha > n$  imply  $\alpha + \beta + \gamma > 3n/2$  and hence it follows from (32) that  $1/p + 1/q + 1/r < 3/2$ . Therefore the plane given by the first equation in (32)



cuts the unit cube  $[0, 1]^3$  at the six points  $A_1 = (1, (2n - \alpha - \beta - \gamma)/n, 0)$ ,  $A_2 = ((2n - \alpha - \beta - \gamma)/n, 1, 0)$ ,  $A_3 = (0, 1, (2n - \alpha - \beta - \gamma)/n)$ ,  $A_4 = (0, (2n - \alpha - \beta - \gamma)/n, 1)$ ,  $A_5 = ((2n - \alpha - \beta - \gamma)/n, 0, 1)$ , and  $A_6 = (1, 0, (2n - \alpha - \beta - \gamma)/n)$ . These six points form the vertices of a hexagon. It suffices to prove Lorentz space estimates at these vertices for characteristic functions. For instance at the vertex  $A_1$  the estimate we need to establish is

$$(33) \quad \int_{E_1} \int_{E_2} \int_{E_3} |x - y|^{-\alpha} |y - z|^{-\beta} |z - x|^{-\gamma} dx dy dz \leq C |E_1| |E_2|^{2 - \frac{\alpha + \beta + \gamma}{n}}.$$

First integrate in  $z$ . We have

$$(34) \quad \int_{E_3} |y - z|^{-\beta} |z - x|^{-\gamma} dz \leq \int_{\mathbf{R}^n} |y - z|^{-\beta} |z - x|^{-\gamma} dz = C |x - y|^{n - \beta - \gamma},$$

for all  $x \neq y$  since  $\beta + \gamma > n$ . The last equality above can be easily shown by a translation, a dilation, and a rotation. Using (34) we obtain

$$\begin{aligned} & \int_{E_1} \int_{E_2} \int_{E_3} |x - y|^{-\alpha} |y - z|^{-\beta} |z - x|^{-\gamma} dx dy dz \\ & \leq C \int_{E_1} \int_{E_2} |x - y|^{n - \alpha - \beta - \gamma} dy dx \\ & \leq C \int_{E_1} \int_{|y| \leq c|E_2|^{1/n}} |y|^{n - \alpha - \beta - \gamma} dy dx \\ & \leq C |E_1| |E_2|^{(2n - \alpha - \beta - \gamma)/n}, \end{aligned}$$

which proves the required estimate (33). This example can be found in [3] when  $n = 1$  and  $\alpha = \beta = \gamma$ .

In this example we have a trilinear form and it is appropriate to apply Corollary 3.8. Again simplifying our conditions on the Boyd indices gives:


**Theorem 5.4.** *Suppose  $X_1, X_2, X_3$  are r.i. spaces on  $\mathbf{R}^n$ . Suppose the Boyd indices satisfy the conditions  $1 < p_{X_i} \leq q_{X_i} < \infty$  for  $i = 1, 2, 3$  and*

$$\sum_{i=1}^3 \frac{1}{q_{X_i}} \leq 3 - \frac{\alpha + \beta + \gamma}{n} \leq \sum_{i=1}^3 \frac{1}{p_{X_i}}.$$

*Then  $I_{\alpha, \beta, \gamma}$  is bounded on  $X_1 \times X_2 \times X_3$  provided the trilinear form  $(f, g, h) \rightarrow x^{2 - \frac{\alpha + \beta + \gamma}{n}} f(x)g(x)h(x)$  is bounded on  $X_1(0, \infty) \times X_2(0, \infty) \times X_3(0, \infty)$ .*

*Remark.* Here as in the preceding example we can relax the conditions on the Boyd indices with the right extra hypotheses. For example if  $p_{X_1} \leq 1$  it is necessary that

$$\frac{1}{q_{X_2}} + \frac{1}{q_{X_3}} > 2 - \frac{\alpha + \beta + \gamma}{n}.$$



**Fig. 1.** The set of all  $(1/p, 1/q, 1/r)$  such that  $|I_{\alpha,\beta,\gamma}(f, g, h)| \leq C \|f\|_{L_p} \|g\|_{L_q} \|h\|_{L_r}$ .

*Example 5.5.* Consider the operator

$$I(f, g)(x) = \int_{|t| \leq 1} f(x + t)g(x - t) dt.$$

We will show  $I$  maps  $L_p(\mathbf{R}^n) \times L_q(\mathbf{R}^n)$  into  $L_r(\mathbf{R}^n)$  when  $(1/p, 1/q, 1/r)$  lies in the closed convex hull of the points  $(1, 0, 0)$ ,  $(0, 1, 0)$ ,  $(1, 0, 1)$ ,  $(0, 1, 1)$ ,  $(1, 1, 1)$ , and  $(1, 1, 1/2)$ .

By interpolation it suffices to establish boundedness estimates at these six points. Five of these estimates are trivial. We only prove that  $I$  maps  $L_1 \times L_1 \rightarrow L_{1/2}$ .

Suppose that we have established the estimate

$$(35) \quad \|I(f, g)\|_{L_{1/2}} \leq C \|f\|_{L_1} \|g\|_{L_1}$$

for all  $f$  and  $g$  supported in two cubes of sidelength one. Then we prove (35) (with a larger constant) for all  $f$  and  $g$  integrable.

For each  $k \in \mathbf{Z}^n$ , let  $Q_k$  be the cube of sidelength one whose sides are parallel to the axes and whose lower left corner is  $k \in \mathbf{Z}^n$ . let  $f_k = f \chi_{Q_k}$  and  $g_m = g \chi_{Q_m}$ . Then for each  $k \in \mathbf{Z}^n$  there exist at most finitely many  $m \in \mathbf{Z}^n$  such that  $I(f_k, g_m)$  is nonzero. This is because the intersection of the sets  $\{t : |t| \leq 1\}$  and  $\frac{1}{2}(Q_k - Q_m)$  has to be nonempty.

Now write

$$I(f, g) = \sum_{k \in \mathbf{Z}^n} \sum_{m \in \mathbf{Z}^n} I(f_k, g_m)$$

as a sum of a finite number of terms of the form

$$\sum_{k \in \mathbf{Z}^n} I(f_k, g_{k+d})$$

where  $d \in \mathbf{Z}^n$  lies in a ball of radius at most a dimensional constant. Now

$$\begin{aligned} \|I(f, g)\|_{L_{1/2}} &\leq \left( \sum_{k \in \mathbf{Z}^n} \int_{\mathbf{R}^n} |I(f_k, g_{k+d})|^{1/2} dx \right)^2 \\ &\leq C \left( \sum_{k \in \mathbf{Z}^n} \|f_k\|_{L_1}^{1/2} \|g_{k+d}\|_{L_1}^{1/2} \right)^2 \leq C \|f\|_{L_1} \|g\|_{L_1}, \end{aligned}$$

by Cauchy-Schwarz, where the penultimate inequality above follows from the assumption that (35) holds for the functions  $f_k$  and  $g_m$ . Summing over  $d$  we obtain the required estimate  $I : L_1 \times L_1 \rightarrow L_{1/2}$  with a larger constant.

We now prove (35) for  $f$  and  $g$  supported in cubes of sidelength one. (Think of  $f = f_k$  and  $g = g_{k+d}$ .) Now observe that  $I(f, g)$  is supported in a cube of sidelength two. Hölder’s inequality gives

$$\begin{aligned} \|I(f, g)\|_{L_{1/2}} &\leq C \|I(f, g)\|_{L_1} \\ &\leq C \int_{\mathbf{R}^n} \int_{|t| \leq 1} |f(x+t)| |g(x-t)| dt dx \leq C \|f\|_{L_1} \|g\|_{L_1}. \end{aligned}$$

*Example 5.6.* We now consider the bilinear fractional integral

$$I_\alpha(f, g)(x) = \int_{\mathbf{R}^n} f(x+t)g(x-t)|t|^{\alpha-n} dt,$$


where  $0 < \alpha < n$ . Homogeneity considerations imply that  $I_\alpha$  can map  $L_p(\mathbf{R}^n) \times L_q(\mathbf{R}^n) \rightarrow L_r(\mathbf{R}^n)$  only when

$$\frac{1}{p} + \frac{1}{q} = \frac{1}{r} + \frac{\alpha}{n}.$$

We will now show that  $I_\alpha$  maps  $L_p \times L_q \rightarrow L_r$  when the point  $(1/p, 1/q, 1/r)$  lies in the open convex hull of the pentagon with vertices  $(\frac{\alpha}{n}, 0, 0)$ ,  $(1, 0, 1 - \frac{\alpha}{n})$ ,  $(1, 1, \frac{2n-\alpha}{n})$ ,  $(0, 1, 1 - \frac{\alpha}{n})$ , and  $(0, \frac{\alpha}{n}, 0)$ . More precisely we will show that a weak-type estimate holds at each vertex of the pentagon below.

We first consider the vertex  $(\frac{\alpha}{n}, 0, 0)$ . Take  $f = \chi_A$  and  $g = \chi_B$ , where  $A$  and  $B$  are measurable sets of finite measure. We have

$$\|I_\alpha(\chi_A, \chi_B)\|_{L_\infty} \leq \sup_{x \in \mathbf{R}^n} \int_{-x+A} |t|^{\alpha-n} dt \leq \int_{|t| \leq c|A|} |t|^{\alpha-n} dt = C|A|^{\alpha/n}.$$



**Fig. 2.** The set of all  $(1/p, 1/q, 1/r)$  such that  $I_\alpha : L_p \times L_q \rightarrow L_r$ .

Likewise we obtain the required estimate at the vertex  $(0, \frac{\alpha}{n}, 0)$ .

The estimates at the vertices  $(1, 0, 1 - \frac{\alpha}{n})$  and  $(0, 1, 1 - \frac{\alpha}{n})$  follow from the estimates at the vertices  $(\frac{\alpha}{n}, 0, 0)$  and  $(0, \frac{\alpha}{n}, 0)$  respectively via duality. Alternatively, just observe that  $I_\alpha(\chi_A, \chi_B) \leq J_\alpha(\chi_A)$ , where  $J_\alpha$  is the usual fractional integral

$$(J_\alpha f)(x) = \int_{\mathbf{R}^n} f(x - y)|y|^{\alpha-n} dy,$$

and thus the estimate  $\|I_\alpha(\chi_A, \chi_B)\|_{L_{n/(n-\alpha),\infty}} \leq C|A|$  directly follows from the corresponding estimate for the linear operator.

Finally we are left with the estimate at the vertex  $(1, 1, \frac{2n-\alpha}{n})$ . For  $j \in \mathbf{Z}$  we introduce operators

$$I_j(f, g)(x) = \int_{|t| \leq 2^j} f(x + t)g(x - t)dt$$

and we note that for  $f, g \geq 0$  we have

$$I_\alpha(f, g) \leq C \sum_{j \in \mathbf{Z}} 2^{j(\alpha-n)} I_j(f, g).$$

Next we observe that by a easy dilation argument  $I_j$  maps  $L_1 \times L_1 \rightarrow L_{1/2}$  with norm bounded by a constant times  $2^{jn}$ . This fact together with the observation

$$\begin{aligned} \int_E (I_j(f, g)(x))^{1/2} dx &\leq \left( \int_E I_j(f, g)(x) dx \right)^{1/2} |E|^{1/2} \\ &\leq C \|f\|_{L_1}^{1/2} \|g\|_{L_1}^{1/2} |E|^{1/2}, \end{aligned}$$

implies that for any measurable set  $E$  with finite measure we have

$$(36) \quad \int_E (I_j(f, g)(x))^{1/2} dx \leq \|f\|_{L_1}^{1/2} \|g\|_{L_1}^{1/2} \min(2^{jn}, |E|)^{1/2}.$$

Now pick  $E = E_\lambda = \{x : |I_\alpha(f, g)(x)| > \lambda\}$ . Then Chebychev’s inequality and (36) give

$$\begin{aligned} \lambda^{1/2} |E_\lambda|^{1/2} &\leq \int_{E_\lambda} \left| \sum_{j \in \mathbf{Z}} 2^{j(\alpha-n)} I_j(f, g)(x) \right|^{1/2} dx \\ &\leq \sum_{j \in \mathbf{Z}} 2^{j(\alpha-n)/2} \int_{E_\lambda} |I_j(f, g)(x)|^{1/2} dx \\ &\leq \sum_{j \in \mathbf{Z}} 2^{j(\alpha-n)/2} \|f\|_{L_1}^{1/2} \|g\|_{L_1}^{1/2} \min(2^{jn}, |E_\lambda|)^{1/2} \\ &= C \|f\|_{L_1}^{1/2} \|g\|_{L_1}^{1/2} |E_\lambda|^{(\alpha-n)/2n}. \end{aligned}$$

This implies that

$$\lambda |E_\lambda|^{\frac{2n-\alpha}{n}} \leq C \|f\|_{L_1} \|g\|_{L_1}$$

which is the required weak type estimate at the vertex  $(1, 1, \frac{2n-\alpha}{n})$ . This example was studied in [4] when  $r \geq 1$  and should be contrasted with the main result in [11]. The same result was independently obtained in [10]. To use the full strength of our results we apply Theorem 4.2 and the succeeding remark to obtain the following generalization for r.i. spaces.

**Theorem 5.7.** *Suppose  $X, Y, Z$  are r.i. spaces on  $\mathbf{R}^n$  with  $Z$  maximal and  $s$ -convex for some  $s > 0$ . Suppose the Boyd indices of  $X, Y, Z$  satisfy the condition that the Boyd cube intersects the pentagon generated by  $(\frac{\alpha}{n}, 0, 0), (1, 0, 1 - \frac{\alpha}{n}), (1, 1, \frac{2n-\alpha}{n}), (0, 1, 1 - \frac{\alpha}{n})$  and  $(0, \frac{\alpha}{n}, 0)$  in a nonempty subset of the interior. Then in order that  $I_\alpha$  maps  $X \times Y$  to  $Z$  it is sufficient that  $(f, g) \rightarrow x^\alpha f(x)g(x)$  maps  $X(0, \infty) \times Y(0, \infty)$  to  $Z(0, \infty)$ .*

## References

1. C. Bennett and R. Sharpley, Interpolation of operators, Pure and Applied Mathematics **129**, Academic Press, Orlando FL, 1988.
2. D. W. Boyd, The Hilbert transform on rearrangement-invariant spaces, *Canad. J. Math.* **19** (1967), 599–616.
3. M. Christ, On the restriction of the Fourier transform to curves: Endpoint results and the degenerate case, *Trans. Amer. Math. Soc.* **287** (1985), 223–238.
4. L. Grafakos, On multilinear fractional integrals, *Studia Math.* **102** (1992), 49–56.
5. S. Janson, On interpolation of multilinear operators, Function spaces and applications (Lund 1986), 290–302, Lecture Notes in Math, **1302**, Springer, Berlin-New York 1988.
6. W. B. Johnson and G. Schechtman, Sums of independent random variables in rearrangement invariant function spaces, *Ann. Probab.* **17** (1989), 789–808.
7. N. J. Kalton, Convexity conditions on non-locally convex lattices, *Glasgow J. Math.* **25** (1984), 141–152.
8. N. J. Kalton, Endomorphisms of symmetric function spaces, *Indiana Univ. Math. J.* **34** (1985), 225–247.
9. N. J. Kalton, J. W. Roberts, and N. T. Peck, An  $F$ -space sampler, *Lon. Math. Soc. Lec. Note Series*, 89, Cambridge University Press, Cambridge-New York, 1984.
10. C. Kenig and E. M. Stein, Multilinear estimates and fractional integration, *Math. Res. Lett.* **6** (1999), 1–15.
11. M. T. Lacey and C.M. Thiele,  $L^p$  bounds for the bilinear Hilbert transform,  $p > 2$ , *Ann. Math.* **146** (1997), 693–724.
12. J. Lindenstrauss and L. Tzafriri, Classical Banach spaces. II. Function spaces, *Ergebnisse der Mathematik und ihrer Grenzgebiete 97*. Springer-Verlag, Berlin-New York, 1979.
13. G. G. Lorentz, Majorants in spaces of integrable functions, *Amer. J. Math.* **77** (1955), 484–492.
14. S. Rolewicz, Metric linear spaces, Second edition, Mathematics and its Applications (East European Series), 20. D. Reidel Publishing Co., Dordrecht-Boston, Mass.; PWN—Polish Scientific Publishers, Warsaw, 1985.
15. R. Sharpley, Interpolation of  $n$  pairs and counterexamples employing indices, *J. Approx. Theory* **13** (1075), 117–127.
16. R. Sharpley, Multilinear weak type interpolation of  $mn$ -tuples with applications, *Studia Math.* **60** (1977), 179–194.
17. R. Strichartz, A multilinear version of the Marcinkiewicz interpolation theorem, *Proc. Amer. Math. Soc.* **21** (1969), 441–444.
18. T. Shimogaki, Hardy-Littlewood majorants in function spaces, *Proc. Japan Acad.* **17** (1965), 365–373.
19. E. M. Stein and G. Weiss, An extension of a theorem of Marcinkiewicz and some of its applications, *J. Math. Mech.* **8** (1959), 263–284.
20. M. Zafran, A multilinear interpolation theorem, *Studia Math.* **62** (1978), 107–124.